US006124458A

United States Patent [19]

Müellen et al.

[11] **Patent Number: 6,124,458**

[45] **Date of Patent: Sep. 26, 2000**

[54] **SUBSTITUTED QUATERRYLENE TETRACARBOXYLIC ACID DIIMIDES**

[75] Inventors: **Klaus Müellen**; **Heribert Quante**, both of Köln; **Arno Böhm**, Mannheim, all of Germany

[73] Assignees: **BASF Aktiengesellschaft**, Ludwigshafen; **Max-Planck-Gesellschaft zur Foerderung der Wissenschaften e.V. Berlin**, Munich, both of Germany

[21] Appl. No.: **09/335,535**

[22] Filed: **Jun. 18, 1999**

Related U.S. Application Data

[62] Division of application No. 08/860,928, filed as application No. PCT/EP90/00118, Jul. 21, 1997, Pat. No. 5,986,099.

[30] **Foreign Application Priority Data**

Jan. 20, 1995 [DE] Germany 195 01 576

[51] **Int. Cl.**[7] **C07D 221/18**

[52] **U.S. Cl.** **546/38**

[58] **Field of Search** 546/38

[56] **References Cited**

U.S. PATENT DOCUMENTS

1,715,430	6/1929	Schmidt et al. .	
4,846,892	7/1989	Henning et al.	106/478
5,405,962	4/1995	Muellen et al.	546/27

FOREIGN PATENT DOCUMENTS

42 36 885 5/1994 Germany .

OTHER PUBLICATIONS

Geerts, et. al., Quaterrylenebis (dicarboximide)s: near infrared absorbing and emitting dyes, Journal of Material Chemistry, vol. 8 (11), 2357–2369, 1998.

Y. Nagao et al, Synthesis and Properties of N–alkylbromoperylene–3,4–dicarboximides, *Dyes and Pigments,* vol. 16, No. 1, Jan. 1, 1991, pp. 19–25, XP000208007.

H. Quante et al, "Quaterrylenebis(dicarboximides)", *Angewandte Chemie International Edition,* vol. 34, No. 12, Jul. 7, 1995, pp. 1323–1325, Weinheim, Germany, XP002002793.

G. Seybold et al, "New Perylene and Violanthrone Dyestuffs for Fluorescent Collectors", *Dyes and Pigments,* vol. 11, No. 4, Jan. 1, 1989, pp. 303–317, XP000084462.

Primary Examiner—Zinna Northington Davis
Assistant Examiner—Binita Robinson
Attorney, Agent, or Firm—Oblon, Spivak, McClelland, Maier & Neustadt, P.C.

[57] **ABSTRACT**

Quaterrylenetetracarboxylic diimides I



I

where

R is hydrogen;

C_1–C_{30}-alkyl whose carbon chain may be interrupted by one or more of —O—, —S—, —NR^1—, —CO— and/or —SO_2— and which may be monosubstituted or polysubstituted by cyano, C_1–C_6-alkoxy or a 5-, 6- or 7-membered heterocyclic radical which is attached via a nitrogen atom and which may contain further heteroatoms and may be aromatic, where

R^1 is hydrogen or C_1–C_6-alkyl;

C_5–C_8-cycloalkyl whose carbon skeleton may be interrupted by one or more of —O—, —S— and/or —NR^1—;

aryl or hetaryl, which may each be monosubstituted or poly-substituted by C_1–C_{18}-alkyl, C_1–C_6-alkoxy, cyano, —$CONHR^2$, —$NHCOR^2$ and/or aryl- or hetaryl-azo, which may each be substituted by C_1–C_{10}-alkyl, C_1–C_6-alkoxy or cyano, where

R^2 is hydrogen; C_1–C_{18}-alkyl; aryl or hetaryl, which may each be substituted by C_1–C_6-alkyl, C_1–C_6-alkoxy, halogen or cyano;

X is halogen; C_1–C_{18}-alkyl; aryloxy, arylthio, hetaryloxy or hetarylthio, which may each be substituted by C_1–C_4-alkyl or C_1–C_4-alkoxy;

n is from 2 to 12,

their preparation and use as pigments or fluorescent dyes and also substituted 9-haloperylene-3,4-dicarbimides III as intermediates therefor.

1 Claim, No Drawings